

**Purdue University**  
**Purdue e-Pubs**

---

LARS Symposia

Laboratory for Applications of Remote Sensing

---

1-1-1977

# On Determining Unharvested Winter Wheat Acreage from Landsat

J. C. Harlan

W. D. Rosenthal

Follow this and additional works at: [http://docs.lib.purdue.edu/lars\\_symp](http://docs.lib.purdue.edu/lars_symp)

---

Harlan, J. C. and Rosenthal, W. D., "On Determining Unharvested Winter Wheat Acreage from Landsat" (1977). *LARS Symposia*. Paper 194.

[http://docs.lib.purdue.edu/lars\\_symp/194](http://docs.lib.purdue.edu/lars_symp/194)

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

Reprinted from

**Symposium on**

**Machine Processing of**

**Remotely Sensed Data**

**June 21 - 23, 1977**

The Laboratory for Applications of  
Remote Sensing

Purdue University  
West Lafayette  
Indiana

IEEE Catalog No.  
77CH1218-7 MPRSD

Copyright © 1977 IEEE  
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to [pubs-permissions@ieee.org](mailto:pubs-permissions@ieee.org).

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

## ON DETERMINING UNHARVESTED WINTER WHEAT ACREAGE FROM LANDSAT

J. C. HARLAN AND W. D. ROSENTHAL

Remote Sensing Center  
Texas A&M University  
College Station, Texas 77843

Nine Landsat passes for one locality and one crop year were examined to determine methods of discriminating harvested from unharvested winter wheat. The nine passes included all growth stages of importance to the Large Area Crop Inventory Experiment (LACIE) including pre-planting and post-harvesting.

Two approaches were utilized in the study. One was based on the examination of the temporal-spectral characteristics of various quality wheat fields within the test site. The other was based on a Landsat wheat yield prediction technique.

Approach 1 requires the examination of the identified wheat fields on each pass with a harvestable/unharvestable decision made for each field after each pass or after some number of passes. The decision can be made on the basis of the change in MSS band radiance curves. This method assumes that all wheat fields can be identified at some point in the crop growth period.

Approach 2 required establishment of a yield estimation technique from Landsat data alone. Based on knowledge of the estimated yield and reasons for abandonment of fields a yield threshold can be set, below which fields will be abandoned (i.e., not harvested).

The following results were obtained:

- MSS band evaluation -  
Each MSS band from 4 to 7 had a lower radiance value when an unharvestable or plowed field was observed. For harvestable fields, the higher the ultimate yield, the greater the difference between band 5 and 6 with 6 being higher. The higher the yield, the more cases where band 7 was higher than band 6.
- Yield -

The yield prediction model was optimized for low yield fields. The model utilized TVI values for fall establishment and tillering, the square of those terms and the value at heading. The model had an  $R^2$  greater than .86 and the variables were shown to be independent.